

DHS FILE COPY

2

AD-A227 030

OFFICE OF NAVAL RESEARCH

Grant N00014-90-J-1193

TECHNICAL REPORT No. 23

Photodissociation Near a Rough Metal Surface: Effect of Reaction Fields

by

Purna C. Das, Ashok Puri and Thomas F. George

Prepared for Publication

in

Journal of Chemical Physics

Departments of Chemistry and Physics
State University of New York at Buffalo
Buffalo, New York 14260

September 1990

Reproduction in whole or in part is permitted for any purpose of the
United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

DTIC
ELECTE
SEP 20 1990
S
E
D
Co

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188


1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/90/TR-23		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York	6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260		7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research	8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Grant N00014-90-J-1193	
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO.	PROJECT NO.
		TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Photodissociation Near a Rough Metal Surface: Effect of Reaction Fields			
12. PERSONAL AUTHOR(S) Purna C. Das, Ashok Puri and Thomas F. George			
13a. TYPE OF REPORT	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) September 1990	15. PAGE COUNT 31
16. SUPPLEMENTARY NOTATION Prepared for publication in Journal of Chemical Physics			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
		PHOTODISSOCIATION, Chemical Reaction, SURFACE PLASMONS	
		ROUGH METAL SURFACE, SILVER SUBSTRATE	
		REACTION FIELDS, SURFACE BUMP. (JS)	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The modification of the photochemical dissociation rate of molecules in the presence of a rough metal surface is explored. Classical electromagnetic calculations are presented for the photodissociation rate of a point dipole near a rough surface modeled as a hemispheroidal bump on a semi-infinite flat plane. A correction is introduced by accounting for the reaction fields due to the dipole-substrate system radiating photons and coupling to delocalized surface plasmons. The effect of the shape and size of the bump and the separation of the molecule from the bump on the rate of photodissociation of the molecule are studied numerically.			
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson		22b. TELEPHONE (Include Area Code) (202) 696-4410	22c. OFFICE SYMBOL

Photodissociation Near a Rough Metal Surface: Effect of Reaction Fields

Purna C. Das
Division of Science, Engineering and Technology
Penn State Erie
The Behrend College
Erie, Pennsylvania 16563-0203

Ashok Puri
Physics Department
University of New Orleans, Lakefront
New Orleans, Louisiana 70148

Thomas F. George
Departments of Chemistry and Physics & Astronomy
Center for Electronic and Electro-optic Materials
239 Fronczak Hall
State University of New York at Buffalo
Buffalo, New York 14260



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Abstract

The modification of the photochemical dissociation rate of molecules in the presence of a rough metal surface is explored. Classical electromagnetic calculations are presented for the photodissociation rate of a point dipole near a rough surface modeled as a hemispheroidal bump on a semi-infinite flat plane. A correction is introduced by accounting for the reaction fields due to the dipole-substrate system radiating photons and coupling to delocalized surface plasmons. The effects of the shape and size of the bump and the separation of the molecule from the bump on the rate of photodissociation of the molecule are studied numerically.

I. Introduction

Ever since the discovery of surface-enhanced Raman scattering (SERS),¹ theoretical and experimental studies have been geared toward understanding the mechanism and possible applications of the interaction of laser radiation with atoms and molecules adsorbed on or near solid surfaces.² Gaining insight into the perturbations induced by the surface is the first step toward spectroscopic applications of a variety of surface optical phenomena which include vibrational spectroscopy of adsorbed molecules by SERS,³ monitoring of surface processes by surface-enhanced nonlinear optics,⁴⁻⁶ possible cooperative phenomena for an ensemble of admolecules,⁷ surface-enhanced fluorescence for fabrication of sensor materials,⁸⁻¹⁰ and possible enhanced photodissociation of adsorbates¹¹⁻¹⁸ with applications in chemical vapor deposition,¹⁹ semiconductor fabrication,²⁰⁻²⁵ and heterogeneous catalysis.^{23,26}

Our interest in this paper lies in the possible enhanced photochemistry of adsorbates. Its potential technological applications to the fabrication and processing of materials depends not only on the cross section for a photodissociation process, but also on the subsequent photodynamics.²⁷ While the photochemistry of gas-phase molecules is relatively well understood, the role of the substrate and the degree of surface roughness in adsorbate photochemistry is a more recent topic of research. The presence of the surface allows energy and electron transfer mechanisms which may inhibit or enhance photofragmentation.¹¹⁻¹⁸ The type of substrate, whether metallic, insulator or semiconductor, also plays an important role in adsorbate photodissociation. Several groups have used insulator and semiconductor substrates, such as LiF(001),¹⁴ Al₂O₃,¹⁸ and Si(111) 7 × 7,²⁵ to study photochemistry. These substrates are good for

detecting and measuring photodissociation and the associated dynamics of fragments, because the nonradiative relaxation rate of molecules on insulator and semiconductor surfaces is much less than that for metals, as predicted by the classical electromagnetic theory.^{28,2,29} Rapid deexcitation of an adsorbate via nonradiative energy transfer to the metallic substrate has been attributed to the absence of photofragmentation of CH_2I_2 on Al or Ag, and phenyloctatetrene on Ag.¹⁸ More recently, for the adsorbate-metal systems $\text{CH}_2\text{CO}/\text{Pt}(111)$ and $\text{CH}_3\text{Br}/\text{Pt}(111)$,¹⁵ and $\text{CH}_3\text{Br}/\text{Ni}(111)$,¹⁶ there is conclusive evidence that photofragmentation does take place without significant surface damping, but the photodynamics is perturbed due to charge transfer and collisional processes.^{17,27} There is also evidence that much of UV surface photochemistry may be charge-transfer mediated at low coverages.^{16,30}

The fragmentation yield is therefore determined by the relative rates of dissociation (direct as well as charge-transfer mediated) and nonradiative energy deposition into the substrate. Since nonradiative energy transfer to,^{28,29} and photoemission from,³¹ a rough surface are radically altered, one expects additional quenching of the photodissociation rate near a rough-surface. Furthermore, a model rough surface may consist of a small particle above a flat surface. Therefore, one may expect rough surface effects to be a combination of the effects of a small particle and the flat plane.³²

While model calculations of a dipole adsorbed on a silver sphere predicts some enhancement effects,¹¹ experimental results are mixed as far as enhancement of photodecomposition rate is concerned. For instance, photodegradation of rhodamine 6G shows no enhancement effects whatsoever.¹² On the other hand, enhanced photodissociation of surface-supported organic

4

molecules has been reported.¹³ The decomposition rate for about four monolayers of pyridine on a roughened silver surface has been found to have a peak for a spacer thickness of ~ 15 to 20 Å. It is believed that the discrepancies between the experiments are a result of the sensitive dependence of the effect on the nature of the surface and on the distance between the molecule and the surface. When chemical decomposition, which follows molecular excitation, is not fast relative to the surface-induced damping processes, which are dominant close to the surface, enhanced molecular absorption does not necessarily imply enhanced photochemical yield. The molecular energy transferred to the substrate, in turn causes excitation of electronic (surface plasmons) or ionic modes of the surface, or electron-hole pairs in the substrate.^{27,33} If the surface modes are radiative,³⁴ part of the energy is recovered, while nonradiative surface modes act as a sink for the energy.

Typically, the electric field in the vicinity of a rough surface or a small particle, irradiated with an external source of light, is increased by two or three orders of magnitude over that in the absence of the surface. Therefore, under steady-state conditions, the power entering or leaving the molecule is increased by four to six orders of magnitude, although the power dissipated into any particular channel of decay is affected by the nature and characteristics of other channels of power flow. Representing the adsorbate-substrate system by a single "system dipole" above a flat surface and allowing this excited dipole to couple to surface plasmons of the underlying flat surface using the energy-transfer theory of Chance, Prock and Silbey,²⁸ and accounting for the loss of power to photon emission and resistive losses in the substrate medium electromagnetically, it has been shown³⁵ that the rates of decay of the dipole to both photons and surface

plasmons are increased substantially, with increasing size of the bump, while the resistive loss rate remained unchanged. Under such enhanced power flow within the molecule-bump system, one is tempted to account for the electromagnetic reactions associated with the various modes of decay of the excited dipole. In fact, radiation damping alone has been shown,³⁶ in the context of SERS from a molecule near a spheroidal particle, to limit enhancement, particularly in the case of large-sized particles. Our goal in this paper, therefore, is to study the effect of correcting the local electromagnetic field, for reactive effects associated with photon radiation as well as coupling to a delocalized surface plasmon by the "system dipole". Within the domain of classical electromagnetism, the corrected local field is seen to better and more accurately predict the sensitive dependence of the photodissociation rate on molecule-surface separation.

The paper is organized in the following manner. In Sec. II, theoretical calculations of the cross section for photodissociation, the enhanced local field and the relative enhancement of molecular photochemistry near a surface are presented. In Sec. III, we present numerical results for the rate of photodissociation and its dependence on the shape and size of the bump, as well as on the separation between the molecule and the bump for a silver substrate, followed by a discussion.

II. Theory

Consider a molecule adsorbed on a rough surface and irradiated by an incident laser, whose electric field vector is denoted by \vec{E}_0 . Let P be the total power transferred to the system while the molecule undergoes photochemical decomposition. The cross section σ for energy transfer to the molecule is defined as

$$\sigma = \left(\frac{8\pi}{c} \right) \Lambda \quad , \quad (1)$$

where $\Lambda = P/|\vec{E}_0|^2$ and c is the speed of light in vacuum. If we denote the quantum efficiency of the molecule for photochemistry by η , then the cross section for photodissociation of the adsorbate is

$$\sigma_{pc} = \eta\sigma = \eta \left(\frac{8\pi}{c} \right) \Lambda \quad . \quad (2)$$

Defining $\sigma_{pc}^0 = \eta \left(\frac{8\pi}{c} \right) \Lambda^0$ as the surface-free cross section of an isolated molecule, we can obtain the surface enhancement ratio for photochemistry:

$$R_{pc} = \Lambda/\Lambda^0 \quad . \quad (3)$$

In the above formalism, we have assumed that the quantum efficiency η is unaffected by the presence of the surface.

A familiar model for a rough surface^{37,35} is shown in Fig. 1. In terms of the spheroidal coordinates (ξ, η, ϕ) , the hemispheroid surface is characterized by $\xi_0 = a/f$; the surface through the position of the molecule, considered as a polarizable point dipole, is given by $\xi_1 = (a + H)/f$, with the size parameter $f = (a^2 - b^2)^{1/2}$; a and b are the semi-major and semi-minor axes of the hemispheroidal protrusion, respectively. The hemispheroid and the flat plane are considered as one piece with a complex dielectric constant $\epsilon(\omega)$. The incident laser field is taken to be propagating along the interface so that its electric field vector \vec{E}_0 is along the normal to the interface. The molecular dipole $\vec{\mu}$ is oriented along the z -direction. The local electric field at the location of the molecular dipole is obtained from the electric potential Φ_{II} in region II,³⁷

$$\Phi_{II} = \sum_n B_n Q_n(\xi) P_n(\eta) - E_0 f \xi \eta$$

$$+ \frac{\mu}{f} \frac{\partial}{\partial \xi_1} \sum_n \frac{(2n+1)}{f} P_n(\xi_{<}) Q_n(\xi_{>}) P_n(\eta) \quad (4)$$

where P_n and Q_n are the Legendre polynomials of the first and second kind, respectively, and $\xi_{<} = \min(\xi, \xi_1)$ and $\xi_{>} = \max(\xi, \xi_1)$. Thus the local electric field at the location of the dipole is given by

$$E_{loc} = E_0 - \underline{B} \cdot \underline{q} \quad (5)$$

where $q_n = Q'_n(\xi_1)/f$.

A formal solution for the coefficients B_n is obtained in the following form (quantities with single and double bars are vectors and matrices, respectively):^{35,37}

$$\underline{B} = (\underline{T}^{-1} \cdot \underline{S} \cdot \underline{b}) \mu - (\underline{T}^{-1} \cdot \underline{R}) E_0 \quad (6)$$

where

$$T_{jn} = X_{jn} \{ [Q_n(\xi_0) P'_j(\xi_0) - \epsilon^{-1} Q'_n(\xi_0) P_j(\xi_0)]$$

$$+ (-1)^{j+n} \Delta_n [Q_n(\xi_0) P'_j(\xi_0) - Q'_n(\xi_0) P_j(\xi_0)] \} \quad (7)$$

$$X_{jn} = \int_0^1 d\eta P_j(\eta) P_n(\eta) \quad (8)$$

$$\Delta_n = \left[\frac{1+\epsilon}{2} \epsilon^{-1} + (-1)^n \frac{1-\epsilon}{2} \epsilon^{-1} \right] , \quad (9)$$

$$S_{jn} = X_{jn} \{ Q'_n(\xi_1) [\epsilon^{-1} P_j(\xi_0) P'_n(\xi_0) - P'_j(\xi_0) P_n(\xi_0)] \\ + (-1)^{j+n} \Delta_n P'_n(\xi_1) [P_j(\xi_0) Q'_n(\xi_0) - P'_j(\xi_0) Q_n(\xi_0)] \} , \quad (10)$$

$$R_j = f \xi_0 \left\{ \frac{2}{3} \delta_{j1} \epsilon^{-1} - X_{j1} P'_j(\xi_0) \left[1 - \frac{(-1)^j}{\epsilon} \right] \right\} , \quad (11)$$

$$b_n = \frac{2n+1}{f^2} . \quad (12)$$

The system dipole moment is given^{35,37} by $D = \left(\frac{B_1 f^2}{3} \right) + \mu$, and E_{loc} is the required field which polarizes the adsorbed molecule. Writing $B_n = W_n \mu - V_n E_0$, where $\underline{W} = \underline{T}^{-1} \cdot (\underline{S} \cdot \underline{b})$ and $\underline{V} = \underline{T}^{-1} \cdot \underline{R}$, one gets

$$E_{loc} = E_0 - (\underline{W} \cdot \underline{q}) \mu + (\underline{V} \cdot \underline{q}) E_0 . \quad (13)$$

$(\underline{V} \cdot \underline{q}) E_0$ is the field produced by the polarization of the surface by the incident field E_0 in the absence of the molecule, and $-(\underline{W} \cdot \underline{q}) \mu$ is the electric field at the location of the molecule produced by the polarization of the surface by the near field of the molecular dipole. By substituting for B_1 we see that the system dipole consists of the molecular dipole μ , a dipole moment $-V_1 E_0 f^2/3$ induced in the bump by the incident field, and a moment $W_1 f^2 \mu/3$ induced in the bump by the molecule.

We are interested in the sum of all the contributions to the local field including the reaction fields, associated with the coupling of the

system dipole to photons and delocalized surface plasmon. Let us write $E_R = RE_0$ and $E_I = G\mu$, where $R(\omega) = \underline{V} \cdot \underline{q}$ and $G(\omega) = -\underline{W} \cdot \underline{q}$. To lowest order, the local field at any point above the surface consists of the incident field, the fields of photons radiated by both the molecular dipole and the dipole induced in the surface protrusion, and the field caused by the excitation of the surface plasmon by both the molecular dipole and the dipole induced in the bump. Therefore, we can imagine the system to consist of a single dipole of moment D , positioned very close to a flat surface.

If we denote the radiation reaction field by E_{rad} and the reaction field associated with the coupling to the surface plasmon by E_{sp} , then

$$E_{\text{loc}} = E'_0 - \underline{B} \cdot \underline{q} \quad , \quad (14)$$

where $E'_0 = E_0 + E_{\text{rad}} + E_{\text{sp}}$, and to maintain self consistency, we use $\underline{B} = \underline{B}(\mu, E'_0)$ by replacing E_0 by E'_0 in Eq. (6), and write the system dipole $D = D(\mu, E'_0)$. The fields, E_{rad} and E_{sp} , are determined by using the results of Ref. 31 for the power radiated to photons,

$$P_{\text{rad}} = \frac{1}{3} |D|^2 \frac{\omega^4}{c^3} \left[1 + \frac{3}{2} \int_0^{\pi/2} d\theta \frac{\sin^3 \theta \{(\epsilon^2 + 1) \cos^2 \theta + (\epsilon - 1)\}}{(\epsilon - 1) \{(\epsilon + 1) \cos^2 \theta - 1\}} \right] \quad , (15)$$

and the power lost to the excitation of surface plasmon,

$$P_{\text{sp}} = 4\pi\omega |D|^2 \left(\frac{\omega}{c}\right)^3 \frac{(-\epsilon)^3}{(-\epsilon - 1)^{5/2} (1 - \epsilon)} \quad , \quad (16)$$

by the system dipole D , and demanding that the work done by the reaction force on the system dipole be equal to the negative of the corresponding energy lost during a given time. This calculation yields

$$E_{\text{rad}} = \frac{iD}{3} \left(\frac{\omega}{c}\right)^3 \left\{ 1 + \frac{3}{2} \int_0^{\pi/2} d\theta \frac{\sin^3 \theta [(\epsilon^2 + 1) \cos^2 \theta + (\epsilon - 1)]}{(\epsilon - 1)[(\epsilon + 1) \cos^2 \theta - 1]} \right\} \quad (17)$$

and

$$E_{\text{sp}} = 4\pi iD \left(\frac{\omega}{c}\right)^3 \frac{(-\epsilon)^3}{(-\epsilon - 1)^{5/2} (1 - \epsilon)} \quad (18)$$

Near the surface, the equation of motion of the molecular dipole, μ , may be taken to be of the Drude form,

$$\ddot{\mu} + \omega_0^2 \mu + \gamma_0 \dot{\mu} = \alpha_0 \omega_0^2 E_{\text{loc}}(t) \quad (19)$$

where ω_0 and γ_0 are the molecular frequency and decay rate in the bulk, respectively. Considering a steady state solution of Eq. (19) in the form $\mu(t) = \mu(\omega)e^{-i\omega t}$, the self-consistent dipole moment μ of the molecule and the power P absorbed by it in the presence of the corrected local field E_{loc} , are straightforwardly obtained:

$$\mu = \frac{\omega_0^2 \alpha_0 K_1 E_0}{(\tilde{\omega}_0^2 - \omega^2 - i\omega\tilde{\gamma})} \quad (20)$$

and

$$P = \frac{\omega_0^2 \alpha_0^2}{2} \frac{|K_1 E_0|^2 \tilde{\gamma}}{(\tilde{\omega}_0^2 - \omega^2)^2 + \omega^2 \tilde{\gamma}^2} \quad (21)$$

where

$$K_1 = [1 + R(\omega)] / (1 + UV_1 f^2 / 3) \quad (22)$$

$$\tilde{\gamma} = \gamma_0 + \frac{\omega_0^2 \alpha_0}{\omega} \text{Im}[G(\omega) + U(1 + \frac{W_1 f^2}{3}) K_1] \quad (23a)$$

$$\tilde{\omega}_0^2 = \omega_0^2 - \omega_0^2 \alpha_0 \text{Re}[G(\omega) + U(1 + \frac{W_1 f^2}{3}) K_1] \quad (23b)$$

and

$$U = U_{\text{rad}} + U_{\text{sp}} \quad (24)$$

is the factor resulting from the inclusion of reaction fields, $E_{\text{rad}} = U_{\text{rad}} D$ and $E_{\text{sp}} = U_{\text{sp}} D$, in the local electric field. V_n and W_n are the n -th elements of vectors \underline{V} and \underline{W} , respectively.

Inspection of Eqs. (20) and (22) shows that when reaction fields are included, the effect of the driving field is cut down by a factor of $(1 + UV_1 f^2 / 3)^{-1}$. Thus the absorbed power is reduced by the absolute square of the above factor from its value in the absence of reaction fields. Also, we find from Eq. (23) that surface-induced width and level shift have terms proportional to UK_1 and are thus dependent on $[1 + R(\omega)]$ when the reaction fields are accounted for. This is typical of situations where effects of the reaction fields are included.³⁶

The ratio of P to P_0 , the power absorbed by the molecule in the absence of any surface gives a measure of the photochemistry enhancement factor R_{pc} . This ratio can be expressed as

$$R_{pc} = \left(\frac{\tilde{\gamma}}{\gamma_0} \right) \left| \frac{E_{loc}}{E_0} \right|^2, \quad (25)$$

or

$$R_{pc} = \left(\frac{\tilde{\gamma}}{\gamma_0} \right) \frac{|K_1|^2 ((\omega_0^2 - \omega^2)^2 + \omega^2 \gamma_0^2)}{((\tilde{\omega}_0^2 - \omega^2)^2 + \omega^2 \tilde{\gamma}^2)}. \quad (26)$$

If we neglect the small level shift (i.e., $\tilde{\omega}_0^2 \approx \omega_0^2$) and assume a near-resonance condition ($\omega \approx \omega_0$), then

$$R_{pc} \approx \left(\frac{\gamma_0}{\tilde{\gamma}} \right) |K_1|^2. \quad (27)$$

This is a familiar expression for R_{pc} , which contains two competing factors. The magnitude of R_{pc} is determined by the field enhancement factor $|K_1|^2$ and by the relative magnitudes of γ_0 and $\tilde{\gamma}$. $|K_1|^2$ contains the normal Raman enhancement factor $|1+R|^2$ and an add-on factor, $|1+UV_1 f^2/3|^{-2}$, due to the inclusion of reactive effects.

III. Results and Discussion

Here, we consider the photodissociation rate of a molecule held at a distance H from a silver surface. The molecule is characterized by the

parameters ω_0 , γ_0 and α_0 . The rough surface consisting of the semi-infinite flat plane and the hemispheroidal protrusion is made up of silver, whose complex dielectric function $\epsilon(\omega)$ is taken from Ref. 38. To obtain finer adjustments in the neighborhood of resonance frequencies (for both molecular and plasmon resonances), we have numerically fitted the data to obtain the following empirical relation for $\epsilon(\omega)$ in the Drude form,

$$\epsilon(\omega) = \left[4.74 + \frac{0.12}{\omega^2} \right] + 0.24i, \quad (28)$$

where ω is the frequency of the incident light. In what follows, all quantities are expressed in atomic units (a.u.).

Numerical calculations are done for the molecule resonating with both the incident laser frequency ω as well as with the ground-state resonance frequency of the bump-plane system, whose surface geometry gives rise to electromagnetic resonances called "shape resonances"³⁵. One way of characterizing the shape resonance is through the value of the real part of the dielectric constant at the resonance frequency. For $a/b = 2, 3$ and 4 , the ground states correspond to $\text{Re}\epsilon(\omega) = -7.34, -11.3$ and -15.9 , respectively. These data, along with Eq. (28), give the ground-state resonance frequency of the bump, which is $0.1, 0.086$ and 0.076 a.u., for aspect ratios of $2, 3$ and 4 , respectively. A ten-dimensional \underline{B} -vector was chosen for numerical computation. For fixed a/b , a and H , reasonable convergence in R_{pc} was obtained using this \underline{B} vector, with $a \leq 1000$ a.u. and $H \leq 140$ a.u.

Figure 2 shows $|K_1|^2$ (----) and $|1 + R(\omega)|^2$ (—) as a function of the distance H for various aspect ratios of the bump. $|1 + R(\omega)|^2$ is

regarded as the field enhancement factor when $U = 0$ and $|K_1|^2$ is the corresponding factor when $U \neq 0$. For a given H , there is a significant reduction in the magnitude of the field enhancement factor when reaction field contributions are included. For example, for $a/b = 2$, there is at least two orders of magnitude reduction. As the bump becomes more needle-shaped, the value of $|K_1|^2/|1 + R(\omega)|^2$ becomes larger, because of the lightning-rod effect. The relative magnitude of $|K_1|^2$ with respect to the normal Raman enhancement factor $|1 + R(\omega)|^2$ for various aspect ratios and semi-major axes of the bump are given in Table 1.

The enhancement factor for photochemistry, R_{pc} , depends on $|K_1|^2$ and $\tilde{\gamma}$, besides other molecular parameters. Since K_1 is the surface-modified field-enhancement factor (when $U \neq 0$) and since it also occurs in $\tilde{\gamma}$, it is easy to see how the molecular width γ changes as a function of H when reaction fields are included. It turns out that the numerical value of $\text{Im}[U(1 + W_1 f^2/3)K_1]$ is small compared to $\text{Im}(G)$, and consequently $\tilde{\gamma}$ changes slightly with respect to γ unless we consider extremely large-sized ($a > 1000$ Å) bumps. $|K_1|^2$ in the numerator of Eq. (26) is thus the dominant factor in deciding the effects of including reaction fields on photochemistry. On the same ground, since the polarizability α_0 enters the expression for R_{pc} through $\tilde{\gamma}$, the effect on R_{pc} of going from weak ($\alpha_0 = 0.1$) to strong ($\alpha_0 = 10.0$) polarizabilities is expected to be minimal.

Figure 3 shows R_{pc} as a function of H for three different aspect ratios for a fixed size of the bump (fixed a). The solid curves are obtained when $U = 0$, and the dashed curves are for $U \neq 0$. The molecular frequency ω_0 is chosen to coincide with the ground state of the bump for each value of a/b . Since R_{pc} is directly proportional to $|K_1|^2$, these curves mimic those of Fig. 2. The reactive effects are evidenced by the

decrease in R_{pc} by several orders of magnitude when $U \neq 0$ from that when $U = 0$. As one goes away from the surface, the enhancement factor declines for a given eccentricity of the hemispheroid, and appreciable enhancement persists out to large distances.

Figure 4 gives the enhancement ratio R_{pc} for a molecule located at 40 a.u. from the surface as a function of the semi-major axis a of the hemispheroid for various aspect ratios. A complete resonance condition ($\omega_0 \approx \omega \approx \omega_s$, where ω_s is the ground-state resonance frequency of the bump-plane system) is assumed. Other molecular parameters used are $\gamma_0 = 10^{-3}$ a.u. and $\alpha_0 = 10$ a.u. These graphs show the effects of both including (----; $U \neq 0$) and excluding (—; $U = 0$) the reaction fields in the calculation. For extremely small values of a (< 200 a.u.), the effect of including the reaction fields is negligible. On the other hand, for a given shape of the bump (a/b fixed), the enhancement with $U \neq 0$ decreases to significantly lower values than those for $U = 0$ with increasing a . This is understandable since the effects of including the reaction fields are expected to be substantial for large-sized roughness features.³⁶ For $a = 300$ Å, it is seen that the reduction in enhancement can be nearly two orders of magnitude when $U \neq 0$ for a hemispheroid with $a/b = 2$. The magnitude of the attenuation in enhancement decreases with increasing eccentricity of the bump, because one would expect the strongest intrinsic electric fields near the sharpest protrusions. The value of a where R_{pc} peaks may be taken as the size of the bump at which reaction field contributions start to bear significant effect. For an eccentricity of 3, for example, the semi-major axis of the bump has to be roughly ~ 300 a.u. before reactive effects are significantly noticeable.

As mentioned earlier, U consists of two parts: $U = U_{\text{rad}} + U_{\text{sp}}$. U_{rad} corresponds to the inclusion of photon radiation damping, and a nonzero U_{sp} represents the inclusion of reaction in the decay of the system dipole to delocalized surface plasmons, which is the primary relaxation mechanism very close to the surface.^{2,28,29} The effect of only $U_{\text{rad}} \neq 0$ was discussed in Ref. 36 in the context of SERS. To see how a nonzero U_{sp} additionally modifies the enhancement of photochemistry near a rough surface, we show R_{pc} as a function of the semi-major axis a in Fig. 5 for three different situations: $U \neq 0$ (curve a), $U = 0$ (curve b) and $U_{\text{rad}} \neq 0$ but $U_{\text{sp}} = 0$ (curve c). Curve c is in qualitative agreement with previous results³⁶ as evidenced by the attenuation of enhancement for larger-sized bumps. Comparative inspection of the curves in Fig. 5 shows that for photodissociation of a molecule near a rough silver surface, the effect of including the surface plasmon reaction field in the calculation is substantial for both intermediate and larger-sized bumps. It clearly illustrates the importance of taking into account the reaction field corresponding to the excitation of surface plasmons by the system dipole. The peak positions of curves a and c set a lower bound for values of the semi-major axis a at which radiation and surface plasmon dampings, respectively, become important.

The above results lead us to a simple conclusion that enhancement in photochemistry near a rough surface is not only dependent, very sensitively, on the shape and size of the roughness feature, but also the theory should take the radiation as well as the surface plasmon damping into consideration, particularly for large-sized features, for a correct prediction of experimental results. If one wants to produce a predetermined enhancement in photochemistry for a molecule held at a fixed distance from

the surface, the surface morphology has to be designed accordingly. On the other hand, if the surface roughness is specified, one needs to put the molecule at a certain, predetermined, distance away from the surface to get maximum enhancement.

The "system dipole" referred to in this article has a moment which is the sum of the molecular dipole moment and the dipole moment associated with the "single" roughness feature. For the near-resonance condition, where the laser frequency coincides with a resonance of the molecule and the bump, each dipole may have its moment increased by a factor of ~50 to 100. Thus the system dipole in this model calculation is much stronger than gas-phase molecular dipole. Consequently, the coupling of an adsorbed molecule to the delocalized surface electronic excitation of the flat substrate in the case of flat-surface adsorbate photochemistry is much weaker than that in the case of a rough surface. One may therefore neglect the reactive effects associated with radiation and surface plasmon dampings in flat surface photochemistry.

Like the energy-transfer mechanism, charge-transfer processes, to³⁹ and from^{17,30} the surface, are short-range quenching mechanisms. The photochemical process involves excitation of the molecule to a repulsive state and subsequent fragmentation. The repulsive state is attained either by electronic excitation or by dissociative electron attachment,^{17,30} where the surface loses an electron to the adsorbate. In either case, quenching of photochemistry occurs very close to the surface (i.e., at very low coverages and/or when the inert spacer layer between the molecule and surface is absent), since the molecule may return to a nonrepulsive state before fragmentation occurs. This is possible because of the strong overlap between the molecule and the surface at short range, so that the attached

electron as well as the excited electron can tunnel through to the surface. On the other hand, very far from the surface, the excited molecule has sufficient time to undergo fragmentation without appreciable quenching by charge transfer. Because of the very weak molecule-surface overlap and the potential barrier that the surface electron has to overcome (to get through the first few layers of molecules or the spacer layer), electron-transfer-mediated photochemistry is practically absent at large distances. There is hence an optimum distance from the surface,¹⁷ where the cross section for photodissociation is expected to peak.

Although from the above discussion, the characteristics of both energy-transfer and charge-transfer quenching mechanisms are similar, we have made no attempt to include the latter effect here. While the energy dissipation to a rough surface can be easily accounted for by the classical electromagnetic theory,³⁷ detailed quantum theoretical insight into the charge-exchange processes in the presence of a "rough" surface is needed.

In summary, we have considered a correction to the electromagnetic theory of surface-enhanced photochemistry by taking into account the radiation and the surface plasmon reaction fields. We chose a model where decay to delocalized plasmons, besides photon emission, by the system is possible. It is found that while surface plasmon damping can be important for both small as well as large-sized roughness features, the radiation damping has an effect only for very large-sized roughness features. It should be pointed out that the rough surface chosen is only a model surface. Realistic rough surfaces are quite different and difficult to model. Nonetheless, the qualitative conclusions drawn from these results should help guide experimentalists in choosing correct parameters to maximize enhancement. Given the surface preparation techniques known to date, it is

perhaps possible to check the predictions of this calculation experimentally on microlithographically-prepared surfaces.⁴⁰

Acknowledgments

This research was supported in part by the National Science Foundation under Grant CHE-8922288 and the Office of Naval Research. PCD wishes to acknowledge the computational assistance of Mr. Donald Wheaton. We also acknowledge valuable discussions with Prof. J. P. Cowin.

a (a.u.)	$ K_1 ^2 / 1 + R(\omega) ^2$		
	a/b = 2	a/b = 3	a/b = 4
400	0.08	0.2	0.3
600	0.01	0.03	0.07
800	0.002	0.008	0.02

Table 1. The ratio of the field-enhancement factor $|K_1|^2$ (when $U \neq 0$) to $|1 + R(\omega)|^2$ (when $U = 0$) as a function of the semi-major axis a of the hemispheroid for different a/b values.

Figure Captions

1. Geometry of the surface protrusion. The semi-major axis is a , and the semi-minor axis is b . The spheroid surface is $\xi = \xi_0$, and the surface passing through the molecule is $\xi = \xi_1$.
2. $|1 + R(\omega)|^2$ (—) and $|K_1|^2$ (----) as a function of the molecule-surface distance H for three different a/b values: (a) $a/b = 2$; (b) $a/b = 3$; (c) $a/b = 4$. For each case, $a = 800$ a.u., $\alpha_0 = 10$ a.u., and ω_0 is equal to the ground-state resonance frequency of the bump-plane system.
3. Enhancement factor R_{pc} as a function of H for three a/b values: (a) $a/b = 2$, $\omega_0 = 0.1$ a.u.; (b) $a/b = 3$, $\omega_0 = 0.086$ a.u.; (c) $a/b = 4$, $\omega_0 = 0.076$ a.u. The solid curves are for $U = 0$, and the dashed curves are for $U \neq 0$. Here, $a = 800$ a.u., $\alpha_0 = 10$ a.u. and $\gamma_0 = 10^{-3}$ a.u.
4. Enhancement factor R_{pc} as a function of the semi-major axis a of the bump for three different aspect ratios: (a) $a/b = 2$, $\omega_0 = 0.1$ a.u.; (b) $a/b = 3$, $\omega_0 = 0.086$ a.u.; (c) $a/b = 4$, $\omega_0 = 0.076$ a.u. The solid curves are for $U = 0$, and the dashed curves are for $U \neq 0$. Here, $H = 40$ a.u., $\alpha_0 = 10$ a.u. and $\gamma_0 = 10^{-3}$ a.u.
5. Enhancement factor R_{pc} as a function of the semi-major axis a for a fixed $a/b = 3$. Curves (a) and (b) correspond to $U \neq 0$ and $U = 0$, respectively. Curve (c) is obtained when only the photon radiation damping is taken into account ($U_{rad} \neq 0$, $U_{sp} = 0$). Here, $H = 40$ a.u., $\omega_0 = 0.086$ a.u., $\alpha_0 = 10$ a.u. and $\gamma_0 = 10^{-3}$ a.u.

References

1. M. Fleischman, P. J. Hendra and A. J. McQuillan, Chem. Phys. Lett. 26, 163 (1974).
2. M. Moskovits, Rev. Mod. Phys. 57, 783 (1985).
3. F. R. Aussenegg, A. Leitner, and M. E. Lippitsch, Eds., Surface Studies with Lasers (Springer, Berlin, 1983).
4. C. K. Chen, T. F. Heinz, D. Ricard, and Y. R. Shen, Phys. Rev. Lett. 46, 1010 (1981); 48, 478 (1982); Y. R. Shen, J. Vac. Sci. Technol. B 3, 1464 (1985); Th. Rasing, Y. R. Shen, M. W. Kim, P. Valint, Jr. and J. Bock, Phys. Rev. A. 31, 537 (1985).
5. G. Marowsky and R. Steinhoff, Opt. Lett. 13, 707 (1988).
6. E. S. Peterson and C. B. Harris, J. Chem. Phys. 91, 2683 (1989).
7. K. C. Liu and T. F. George, Phys. Rev. B 32, 3622 (1985); Surf. Sci. 164, 149 (1985).
8. D. A. Weitz, S. Garoff, J. I. Gersten and A. Nitzan, J. Chem. Phys. 78, 5324 (1983) and references therein.
9. X. Y. Huang, J. T. Lin and T. F. George, J. Chem. Phys. 80, 893 (1984); X. Y. Huang, K. T. Lee and T. F. George, J. Chem. Phys. 85, 567 (1986).
10. K. C. Lee, Surf. Sci. 163, L759 (1985); M. Lieberherr, Ch. Fattinger and W. Lukosz, Surf. Sci. 189/190, 954 (1987); E. H. Hellen and D. Axelrod, J. Opt. Soc. Am. 4, 337 (1987); F. R. Aussenegg, A. Leitner, M. E. Lippitsch, H. Reinisch and M. Riegler, Surf. Sci. 189/190, 935 (1987).
11. A. Nitzan and L. E. Brus, J. Chem. Phys. 75, 2205 (1981).
12. S. Garoff, D. A. Weitz and M. S. Alvarez, Chem. Phys. Lett. 93, 283 (1982); S. Garoff, D. A. Weitz, M. S. Alvarez and J. I. Gersten, J. Chem. Phys. 81, 5189 (1984); J. I. Gersten and A. Nitzan, Surf. Sci.

- 158, 165 (1985); P. Das and H. Metiu, J. Phys. Chem. 89, 4680 (1985); P. T. Leung and T. F. George, J. Chem. Phys. 85, 4729 (1986).
13. G. M. Goncher and C. B. Harris, J. Phys. Chem. 77, 3767 (1982); P. M. Whitmore, A. P. Alivisatos and C. B. Harris, Phys. Rev. Lett. 50, 1092 (1983); C. J. Chen and R. M. Osgood, Appl. Phys. A 31, 171 (1983); G. M. Goncher, C. A. Parsons and C. B. Harris, J. Phys. Chem. 88, 4200 (1984).
14. F. Tabares, E. P. Marsh, G. A. Bach and J. P. Cowin, J. Chem. Phys. 86, 738 (1987); I. Harrison, J. C. Polanyi and P. A. Young, J. Chem. Phys. 89, 1475, 1498 (1988).
15. S. A. Costello, B. Roop, Z. M. Liu, and J. M. White, J. Phys. Chem. 92, 1019 (1988); B. Roop, S. A. Costello, C. M. Greenlief and J. M. White, Chem. Phys. Lett. 143, 38 (1988).
16. E. P. Marsh, M. R. Schneider, T. L. Gilton, F. L. Tabares, W. Meier and J. P. Cowin, Phys. Rev. Lett. 60, 2551 (1988); E. P. Marsh, F. L. Tabares, M. R. Schneider, T. L. Gilton, W. Meier and J. P. Cowin, J. Chem. Phys. 92, 2004 (1990).
17. E. P. Marsh, T. L. Gilton, W. Meier, M. R. Schneider, and J. P. Cowin, Phys. Rev. Lett. 61, 2725 (1988).
18. T. J. Chuang and K. Domen, J. Vac. Sci. Technol. A 5, 473 (1987); K. Domen and T. J. Chuang, Phys. Rev. Lett. 59, 1484 (1987).
19. S. R. J. Brueck and D. J. Ehrlich, Phys. Rev. Lett. 48, 1678 (1982); F. Bozso and Ph. Avouris, Appl. Phys. Lett. 53, 1095 (1988).
20. F. A. Houle, T. F. Deutsch and R. M. Osgood, Jr., Eds., Laser Chemical Processing of Semiconductor Surfaces, (Materials Research Society, Pittsburgh, 1984).

21. P. D. Brewer, D. McClure and R. M. Osgood, Jr., Appl. Phys. Lett. 49, 803 (1986).
22. J. H. Fendler, J. Phys. Chem. 89, 2730 (1985).
23. T. J. Chuang, Surf. Sci. Rep. 3, 1 (1983).
24. A. P. Alivisatos, M. F. Arndt, S. Efrima, D. H. Waldeck and C. B. Harris, J. Chem. Phys., 86, 6540 (1987); and references therein.
25. C. E. Bartosch, N. S. Gluck, W. Ho and Z. Ying, Phys. Rev. Lett. 57, 1425 (1986).
26. J. H. Snifelt, in Many-Body Phenomena at Surfaces, ed. by D. Langreth and H. Suhl (Academic, Orlando, 1984).
27. Ph. Avouris and B. N. J. Persson, J. Phys. Chem. 88, 837 (1984).
28. R. R. Chance, A. Prock and R. Silbey, Adv. Chem. Phys. 37, 1 (1978).
29. H. Metiu, Prog. Surf. Sci. 77, 153 (1986).
30. M. R. Schneider, C. P. Dehnhostel, T. L. Gilton, and J. P. Cowin (preprint).
31. A. Schmidt-Ott, P. Schurtenberger, and H. C. Siegmann, Phys. Rev. Lett. 45, 1284 (1980); U. Even, K. A. Holcomb, C. W. Snyder, P. R. Antoniewicz, and J. C. Thompson, Surf. Sci. 165, L35 (1986).
32. P. K. Aravind and H. Metiu, Surf. Sci. 124, 124 (1983); R. Rupp, Surf. Sci. 127, 108 (1983); T. Takemori, M. Inoue and K. Ohtaka, J. Phys. Soc. Japan. 56, 1587 (1987).
33. M. Persson and B. Hellsing, Phys. Rev. Lett. 49, 662 (1982); B. N. J. Persson and N. D. Lang, Phys. Rev. B 26, 5409 (1982).
34. G. S. Agarwal and C. V. Kunasz, Phys. Rev. B 26, 5832 (1982).
35. P. C. Das and J. I. Gersten, Phys. Rev. B 25, 6281 (1982).
36. A. Wokaun, J. P. Gordon and P. F. Liao, Phys. Rev. Lett. 48, 957 (1982).

37. J. I. Gersten and A. Nitzan, J. Chem. Phys. 73, 3023 (1980).
38. H. J. Hagemann, W. Gudat and C. Kuntz, DESY Report No. SR 7417 (May 1974).
39. F. Buzso, C. P. Hanrahan, J. Arias, J. T. Yates, Jr., H. Metiu and R. M. Martin, Surf. Sci. Lett. 128, 197 (1983); F. Bozso, J. Arias, C. P. Hanrahan, R. M. Martin, J. T. Yates, Jr. and H. Metiu, Surf. Sci. 136, 257 (1984).
40. P. F. Liao, J. G. Bergman, D. S. Chemla, A. Wokaun, J. Melngailis, A. M. Hawryluk and N. P. Economou, Chem. Phys. Lett. 82, 355 (1981).

Fig. 1




Fig. 2




Fig. 3




Fig. 4





Fig. 5



TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

Office of Naval Research (2)
Chemistry Division, Code 1113
800 North Quincy Street
Arlington, Virginia 22217-5000

Commanding Officer (1)
Naval Weapons Support Center
Dr. Bernard E. Douda
Crane, Indiana 47522-5050

Dr. Richard W. Drisko (1)
Naval Civil Engineering
Laboratory
Code L52
Port Hueneme, CA 93043

David Taylor Research Center (1)
Dr. Eugene C. Fischer
Annapolis, MD 21402-5067

Dr. James S. Murday (1)
Chemistry Division, Code 6100
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. David L. Nelson (1)
Chemistry Division
Office of Naval Research
800 North Quincy Street
Arlington, Virginia 22217

Dr. Robert Green, Director (1)
Chemistry Division, Code 385
Naval Weapons Center
China Lake, CA 93555-6001

Chief of Naval Research (1)
Special Assistant for Marine
Corps Matters
Code 00MC
800 North Quincy Street
Arlington, VA 22217-5000

Dr. Bernadette Eichinger (1)
Naval Ship Systems Engineering
Station
Code 053
Philadelphia Naval Base
Philadelphia, PA 19112

Dr. Sachio Yamamoto (1)
Naval Ocean Systems Center
Code 52
San Diego, CA 92152-5000

Dr. Harold H. Singerman (1)
David Taylor Research Center
Code 283
Annapolis, MD 21402-5067

Defense Technical Information Center (2)
Building 5, Cameron Station
Alexandria, VA 22314

FY90 Abstracts Distribution List for Solid State & Surface Chemistry

Professor John Baldeschwieler
Department of Chemistry
California Inst. of Technology
Pasadena, CA 91125

Professor Paul Barbara
Department of Chemistry
University of Minnesota
Minneapolis, MN 55455-0431

Dr. Duncan Brown
Advanced Technology Materials
520-B Danury Rd.
New Milford, CT 06776

Professor Stanley Bruckenstein
Department of Chemistry
State University of New York
Buffalo, NY 14214

Professor Carolyn Cassady
Department of Chemistry
Miami University
Oxford, OH 45056

Professor R.P.H. Chang
Dept. Matls. Sci. & Engineering
Northwestern University
Evanston, IL 60208

Professor Frank DiSalvo
Department of Chemistry
Cornell University
Ithaca, NY 14853

Dr. James Duncan
Federal Systems Division
Eastman Kodak Company
Rochester, NY 14650-2156

Professor Arthur Ellis
Department of Chemistry
University of Wisconsin
Madison, WI 53706

Professor Mustafa El-Sayed
Department of Chemistry
University of California
Los Angeles, CA 90024

Professor John Eyler
Department of Chemistry
University of Florida
Gainesville, FL 32611

Professor James Garvey
Department of Chemistry
State University of New York
Buffalo, NY 14214

Professor Steven George
Department of Chemistry
Stanford University
Stanford, CA 94305

Professor Tom George
Dept. of Chemistry & Physics
State University of New York
Buffalo, NY 14260

Dr. Robert Hamers
IBM T.J. Watson Research Center
P.O. Box 218
Yorktown Heights, NY 10598

Professor Paul Hansma
Department of Physics
University of California
Santa Barbara, CA 93106

Professor Charles Harris
Department of Chemistry
University of California
Berkeley, CA 94720

Professor John Hemminger
Department of Chemistry
University of California
Irvine, CA 92717

Professor Roald Hoffmann
Department of Chemistry
Cornell University
Ithaca, NY 14853

Professor Leonard Interrante
Department of Chemistry
Rensselaer Polytechnic Institute
Troy, NY 12181

Professor Eugene Irene
Department of Chemistry
University of North Carolina
Chapel Hill, NC 27514

Dr. Sylvia Johnson
SRI International
333 Ravenswood Avenue
Menlo Park, CA 94025

Dr. Zakya Kafafi
Code 6551
Naval Research Laboratory
Washington, DC 20375-5000

Professor Larry Kesmodel
Department of Physics
Indiana University
Bloomington, IN 47403

Professor Max Lagally
Dept. Metal. & Min. Engineering
University of Wisconsin
Madison, WI 53706

Dr. Stephen Lieberman
Code 522
Naval Ocean Systems Center
San Diego, CA 92152

Professor M.C. Lin
Department of Chemistry
Emory University
Atlanta, GA 30322

Professor Fred McLafferty
Department of Chemistry
Cornell University
Ithaca, NY 14853-1301

Professor Horia Metiu
Department of Chemistry
University of California
Santa Barbara, CA 93106

Professor Larry Miller
Department of Chemistry
University of Minnesota
Minneapolis, MN 55455-0431

Professor George Morrison
Department of Chemistry
Cornell University
Ithaca, NY 14853

Professor Daniel Neumark
Department of Chemistry
University of California
Berkeley, CA 94720

Professor David Ramaker
Department of Chemistry
George Washington University
Washington, DC 20052

Dr. Gary Rubloff
IBM T.J. Watson Research Center
P.O. Box 218
Yorktown Heights, NY 10598

Professor Richard Smalley
Department of Chemistry
Rice University
P.O. Box 1892
Houston, TX 77251

Professor Gerald Stringfellow
Dept. of Matls. Sci. & Engineering
University of Utah
Salt Lake City, UT 84112

Professor Galen Stucky
Department of Chemistry
University of California
Santa Barbara, CA 93106

Professor H. Tachikawa
Department of Chemistry
Jackson State University
Jackson, MI 39217-0510

Professor William Unertl
Lab. for Surface Sci. & Technology
University of Maine
Orono, ME 04469

Dr. Terrell Vanderah
Code 3854
Naval Weapons Center
China Lake, CA 93555

Professor John Weaver
Dept. of Chem. & Mat. Sciences
University of Minnesota
Minneapolis, MN 55455

Professor Brad Weiner
Department of Chemistry
University of Puerto Rico
Rio Piedras, Puerto Rico 00931

Professor Robert Whetten
Department of Chemistry
University of California
Los Angeles, CA 90024

Professor R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, CA 90024

Professor Nicholas Winograd
Department of Chemistry
Pennsylvania State University
University Park, PA 16802

Professor Aaron Wold
Department of Chemistry
Brown University
Providence, RI 02912

Professor Vicki Wysocki
Department of Chemistry
Virginia Commonwealth University
Richmond, VA 23284-2006

Professor John Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, PA 15260